

A PARTICLE METHOD WITH GLOBAL SUPERCONVERGENCE

S. Hao^a, W. Han^b

^aDepartment of Mechanical Engineering/Department of Materials Science Engineering
Northwestern University, Evanston, Illinois 60208
suhao@northwestern.edu

^bDepartment of Mathematics
University of Iowa
Iowa City, IA 52242
whan@math.uiowa.edu

A method to gain a global superconvergence has been developed for meshfree methods [1, 2] and for the MPFEM – a finite element-based particle method [3]. We first discuss the definition of “superconvergence”. In general a particle method approximates solution via the relation:

$$u^h(x) = \sum_{i=1}^N F_i(x, y_i) \cdot u_i \quad \text{and} \quad F_i(x, y_i) = \phi(y_i - x) \cdot c(x) \cdot \Delta v_i \quad (1)$$

where $\phi(z)$, $c(x)$, and Δv_i are window function, correction function, and a weight at y_i . The relation (1) can be interpreted as the transformation from a set of samples u_i to the approximated solution u^h through the filter F . The convergence rate of u^h , e.g. L_2 norm, is determined by the functional properties of $c(x)$ and $\phi(y_i - x)$ through the satisfaction of the imposed consistence conditions [4-6]. We assume that $c(x)$ is a m_c order polynomial and the degree of freedom of $\phi(z)$, i.e., the number of the undetermined constants in $\phi(z)$, is m_ϕ . Then, in general the convergence rates of (1) for $u^h(x)$ and its n^{th} derivatives, denoted as “ m ” and “ m_n ”, respectively, are:

$$m = m_c + m_\phi + 1; \quad m_n = m_c + m_\phi + 1 - n$$

For example, when $\phi(z)$ is defined by a 5-order spline function ($m_\phi=0$) and $c(x)$ is a linear function ($m_c=1$), usually the L_2 norm for the meshfree methods with (1) is 2.

Analogue to the superconvergence analysis in finite element [6], we define a particle method with superconvergence when

$$m > m_c + m_\phi + 1 \quad \text{and/or} \quad m_n > m_c + m_\phi + 1 - n$$

In the proposed method, the global superconvergence is achieved by a two-level interpolation scheme that allows the satisfaction of higher order reproducing condition for given m_ϕ and m_c .

References

- [1] T. Belytschko, Y. Lu, and L. Gu, *Int. J. for Num. Methods Eng.*, v. 37, p. 229-256, 1994.
- [2] W. K. Liu, S. Jun, Y. F. Zhang, *Int. J. for Num. Methods in Fluids*, 20: p. 1081-1106, 1995.
- [3] S. Hao, H. S. Park, W. K. Liu, *Int. J. for Num. Methods Eng.*, 53(8): p. 1937-1958, 2002.
- [4] Belytschko, T., et al., *Int. J. for Num. Methods Eng.*, 43(5): p. 785-803, 1998.
- [5] Liu, W.K., Chen, Y.J., Uras, R.A., and Chang, C.T., *Comp. Methods in Appl. Mech. Eng.*, 139(1-4): p. 91-157, 1996.
- [6] Babuska, I., and Melenk, J.M., v. 40, p. 727-758, 1996.
- [7] Wahlbin, L.B., *Superconvergence in Galerkin Finite Element Methods*. Lecture Notes in Mathematics 1605, ed. A.D. Heidelberg and F.T. Groningen. 1995: Springer.